



UNIVERSITI PUTRA MALAYSIA

**EVALUATION OF SPATIAL VARIABILITY OF SOIL AND LEAF
TISSUE NUTRIENT STATUS AND YIELD (FRESH FRUIT BUNCH)
IN AN OIL PALM PLANTATION**

TAMALUDDIN SYAM

FH 2002 22

**EVALUATION OF SPATIAL VARIABILITY OF SOIL AND LEAF TISSUE
NUTRIENT STATUS AND YIELD (FRESH FRUIT BUNCH)
IN AN OIL PALM PLANTATION**

By

TAMALUDDIN SYAM

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia, in Fulfillment of the Requirement
for the Degree of Doctor of Philosophy**

November 2002



DEDICATION

This thesis is dedicated to:

My beloved parents,
my wife, my daughter, my families

Abstract of thesis submitted to the Senate of Universiti Putra Malaysia
in fulfilment of the requirement for the degree of Doctor of Philosophy

**EVALUATION OF SPATIAL VARIABILITY OF SOIL AND LEAF TISSUE
NUTRIENT STATUS AND YIELD (FRESH FRUIT BUNCH)
IN AN OIL PALM PLANTATION**

By

TAMALUDDIN SYAM

November 2002

Chairman: Major Professor Dr. Kamaruzaman Jusoff

Faculty: Faculty of Forestry

The study was carried out at Dusun Durian Estate, Golden Hope Plantations Berhad Banting, Selangor, in a 50 ha plot which represents a coastal oil palm plantation. The general objective of this study was to obtain accurate and timely information about the spatial distribution and status of nutrients in the soil and leaf tissue using Geographic Information System (GIS) and remote sensing technology for precision farming of oil palm plantation. Collection of soil and leaf tissue data were conducted by using systematic sampling and an Omni Star DGPS was used to precisely determine the sample sites location. Geostatistics (GS+) software and classical statistics were used for data analysis. A SPOT image of K-J.299344 acquired on March 2000 from the Malaysian Centre for Remote Sensing (MACRES) was analysed using a PCI software version 7.0. Descriptive statistical analysis classed the status of total N as low, whereas available P and exchangeable K varied from moderate to high. Soil exchangeable Ca and Mg ranged from low to moderate. Soil nutrients variability of available P, exchangeable K, Ca and Mg status were classified as high and moderate for total N. Semivariance analysis showed that the

total N, available P and exchangeable K in the soil have a moderately spatial dependence and while strongly spatial dependence for exchangeable Ca and Mg with available lag distance of 52 to 117 m. Leaf nutrient analysis illustrated that the leaf nutrient variability of N, P, and K could be classified as low (CV values of 7, 8 and 12) and moderate variability for Ca and Mg (CV values of 19 and 22). Based on the semivariance analysis, all the leaf nutrients have a moderately spatial dependence with the lag distances of 75, 75, 51, 63 and 117 m for N, P, K, Ca and Mg, respectively. There was no strong relationship between nutrient content in the soil and nutrient content in leaf tissue; therefore nutrient content analysis in leaf tissue was not able to predict nutrient content in the soil. Yield of fresh fruit bunches (FFB) has a high variability and moderately spatial dependence within the lag distance of 84 m. Both the soil nutrient and nutrient content in leaf tissue were not able to predict FFB, because the statistical analysis indicated that there were no strong relationships between yield FFB and nutrients in soil and leaf tissue. SPOT image analysis using Digital Numbers (DN), unsupervised and supervised classifications methods could not be used to predict nutrient content in leaf tissue.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENILAIAN VARIASI RUANG KANDUNGAN NUTRIEN TANAH
DAN DAUN SERTA HASIL (BUAH TANDAN SEGAR)
PADA LADANG KELAPA SAWIT**

Oleh

TAMALUDDIN SYAM

November 2002

Pengerusi: Mejar Profesor Dr. Kamaruzaman Jusoff

Fakulti : Faculty of Forestry

Satu kajian telah di jalankan di kawasan pesisiran pantai Ladang Dusun Durian, Golden Hope Plantations Berhad, di Banting, Selangor dengan keluasan 50 ha. Secara amnya tujuan penyelidikan ialah untuk memperolehi data yang pasti dan tepat masa hal pergantungan ruang dan kadar nutrien di dalam tanah dan di dalam daun menggunakan sistem maklumat sejagat (GIS) dan penderiaan jauh untuk pertanian tepat pada kelapa sawit. Pengumpulan sampel tanah dan daun dikerjakan secara beraturan dengan bantuan Omni Star DGPS untuk penetapan kedudukan pasti daripada sample. Perisian geostatistics (GS+) dan statistik biasa telahpun digunakan untuk analisis data. Satu imej SPOT bernombor K-J. 299344 liputan Mac 2000 yang diperolehi daripada Pusat Remote Sensing Malaysia (MACRES) telahpun dianalisis menggunakan perisian PCI versi 7.0. Uraian analisis statistik menunjukkan bahawa status total N di dalam tanah telahpun dikelaskan ke dalam tingkatan rendah, sedangkan P boleh di dapati dan K boleh dipertukarkan berbagai-bagai daripada sederhana hingga tinggi. Sementara itu, Ca dan Mg boleh dipertukarkan berbagai-

bagai antara tingkatan rendah hingga tinggi. Variasi nutrien tanah P boleh di dapati, K, Ca dan Mg boleh ditukar telah dikelaskan kepada variasi yang besar, sedangkan untuk total N mempunyai variasi sederhana. Semivarian analisis juga menunjukkan bahawa total N, P boleh didapati dan K boleh ditukar di dalam tanah mempunyai pergantungan ruang sederhana, sedangkan untuk Ca dan Mg boleh ditukar mempunyai pergantungan ruang yang kuat dengan jarak berbagai-bagai daripada 50 m sehingga 117 m. Analisis nutrien di dalam daun menunjukkan bahawa nutrien di dalam daun daripada N, P and K boleh di kelaskan kepada variasi kecil ($CV = 7, 8$ dan 12), manakala untuk nutrien Ca dan Mg dikelaskan kepada variasi sederhana ($CV = 19$ dan 22). Didasarkan kepada semivarian analisis menunjukkan bahawa semua nutrien di dalam daun mempunyai pergantungan ruang sederhana dengan jarak 75, 75, 51, 63 dan 117 m, berturut-turut untuk N, P, K, Ca dan Mg. Tiada hubung kait yang kuat antara nutrien di dalam tanah dengan nutrien di dalam daun, hal ini menyifatkan bahawa kandungan nutrien di dalam daun tidak boleh digunakan untuk meramalkan nutrien yang ada di dalam tanah. Hasil buah tandan segar (FFB) mempunyai variasi yang besar dan kebergantungan ruang yang sederhana dengan jarak 84 m. Kadar nutrien di dalam tanah mahupun nutrien di dalam daun tidak boleh digunakan untuk meramalkan hasil FFB kerana tiada hubungan yang bermakna antara FFB dengan nutrien di dalam tanah ataupun nutrien di dalam daun. Analisis imej SPOT menggunakan kaedah angka digital (DN), pengkelasan terselia dan tanpa selia tidak boleh digunakan untuk meramalkan kadar nutrien di dalam daun.

ACKNOWLEDGEMENTS

First and foremost, I would like to express my sincere gratitude to my major supervisor, Major Professor Dr. Kamaruzaman Jusoff, for his guidance, encouragement, ancillary support and friendship. I am also grateful to Associate Professor Dr. Anuar Abd. Rahim, Dr. Mohd. Hashim Tajudin, Dr. Zin Zawawi Zakaria and Dr. Adzmi Mad Arshad, respectively, the supervisory committee members, for their guidance, valuable suggestions and constructive comments throughout the study.

Special thanks are due to Major Professor Dr. Kamaruzaman Jusoff of the Forest Geospatial Information and Survey Laboratory, Universiti Putra Malaysia (UPM) for providing financial support as Graduate Research Assistanship to undertake this study at UPM. I would also like to thank Dr. Mohd. Hashim Tajudin, Director Research and Development Department, Golden Hope Plantations Berhad, Golden Hope Research Centre and his staff for their full support during the research. My sincere thanks go the SEAMEO SEARCA for providing thesis grant award and also to the Malaysian Centre for Remote Sensing (MACRES) for the SPOT image.

I would also like to express my gratitude and sincere appreciation to all Field Research Assistants especially Pakcik Abdullah, Pakcik Sidek, Cik Ashwan and Zaki, who have rendered services in the conduct of this study. I would like to extend my thanks to my friends and colleagues who have contributed their support directly or indirectly to this study.

Last but not least, I would like to express my deepest appreciation to my parents and family members for their moral support. To my wife Masturo Syarif and my daughter Putri Rachma Tasya who not only endured without protest the loneliness but also provided their loving encouragement throughout this long study.

I certify that an Examination Committee met on 27th November 2002 to conduct the final examination of Tamaluddin Syam on his Doctor of Philosophy thesis entitled "Evaluation of Spatial Variability of Soil and Leaf Tissue Nutrient Status and Yield (Fresh Fruit Bunch) in an Oil Palm Plantation" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Ahmad Ainuddin Nuruddin, Ph.D.

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Chairman)

Kamaruzaman Jusoff, Ph.D.

Professor and Coordinator
Forest Geospatial Information and Survey Laboratory
Faculty of Forestry
Universiti Putra Malaysia
(Member)

Anuar Abd. Rahim, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Adzemi Mat Arshad, Ph.D.

Lecturer
Faculty of Agro-technology and Food Science
Kolej Universiti Sains & Teknologi Malaysia
Terengganu Darul Iman
(Member)

Mohd. Hashim Tajudin, Ph.D.

Director, R & D Department
Golden Hope Plantations Bhd.
Golden Hope Research Centre
(Member)

Zin Zawawi Hj. Zakaria, Ph.D.

Head
Smallholders Development and Transfer of Technology
Malaysian Palm Oil Board (MPOB)
(Member)

Pierre C. Robert, Ph.D.

Professor and Director
Precision Agriculture Centre
University of Minnesota, USA
(Independent Examiner)



SHAMSHER MOHAMAD RAMADILI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **19 FEB 2003**

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Kamaruzaman Jusoff, Ph.D.

Professor and Coordinator
Forest Geospatial Information and Survey Laboratory
Faculty of Forestry
Universiti Putra Malaysia
(Chairman)

Anuar Abd. Rahim, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Adzemi Mat Arshad, Ph.D.

Lecturer
Faculty of Agrotechnology and Food Science
Kolej Universiti Sains & Teknologi Malaysia
Terengganu Darul Iman
(Member)

Mohd. Hashim Tajudin, Ph.D.

Director, R & D Department
Golden Hope Plantations Bhd.
Golden Hope Research Centre
(Member)

Zin Zawawi Hj. Zakaria, Ph.D.

Head
Smallholders Development and Transfer of Technology
Malaysian Palm Oil Board (MPOB)
(Member)

AINI IDERIS, Ph.D.

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



TAMALUDDIN SYAM

Date: 18 February 2003

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
LIST OF TABLES	xiv
LIST OF FIGURES	xvii
 I. INTRODUCTION	 1
Background of the Study	1
Problem Statement	3
Objective of Study	5
 II. LITERATURE REVIEW	 7
Precision Farming	7
Introduction	7
Scope of Precision Farming	8
Strategy of Precision Farming	8
Global Positioning System (GPS) and Precision Farming	19
Global Positioning System Overview	19
Differential Global Positioning System (DGPS)	22
Global Positioning System in Precision Farming	24
Geographic Information System (GIS)	27
What is Geographic Information System	27
Maps and Spatial Information	30
Geographic Information System in Precision Farming	31
Remote Sensing in Precision Farming	45
Application of Remote Sensing and Geographic Information System in Yield Forecasting	55 63
Soil Nutrients	64
Nitrogen (N)	67
Phosphorus (P)	70
Potassium (K)	73
Calcium (Ca)	74
Magnesium (Mg)	76
Nutrients in Leaf Tissue	
 III. MATERIALS AND METHODS	 81
Site Description	81
Site Location	81
Climate	81
Management Practices	83
Annual Production of Fresh Fruit Bunches (FFB)	86
Data Collection	87
Satellite Image	87

Leaf Data	88
Soil Data	89
Fresh Fruit Bunches (FFB)	90
Secondary Data	91
Data Analysis	92
Image Analysis	93
Soil Analysis	97
Leaf Tissue Analysis	100
Differential Global Positioning System Measurement	102
Statistical Analysis	103
Geostatistical Analysis	105
MapInfo Analysis	106
IV. RESULT AND DISCUSSION	107
Spatial Nutrient Variability	107
Spatial Variability of N in Soil and Leaf Tissue	107
Spatial Variability of P in Soil and in Leaf Tissue	120
Spatial Variability of K in Soil and Leaf Tissue	132
Spatial Variability of Ca in Soil and Leaf Tissue	144
Spatial Variability of Mg in Soil and Leaf Tissue	154
Spatial Variability of Fresh Fruit Bunch (FFB)	164
SPOT Image Analysis	170
V. CONCLUSIONS AND RECOMMENDATION	176
Conclusions	176
Recommendation.....	177
REFERENCES	179
APPENDICES	198

LIST OF TABLES

Table	Page
1 Locations of the components of the operation control segment	21
2 Examples of synonymous terms for geographic information system and the source or motivation behind their derivation	28
3 Validity of yield estimates in 1993-1994 Rabi Season.....	56
4 Correlation between yield and NDVI at different stages of paddy crop over CCE plots	57
5 Nutrients removed by various plantation crops	63
6 Effects of N fertilizers on FFB yield at Sg. Wakuba Estate (1990-1996)	66
7 Effects of P fertilizers on FFB yield at Sg. Wakuba Estate (1990-1996)	69
8 Effects of K fertilizers on FFB yield at Sg. Wakuba Estate (1990-1996)	72
9 Yield responses to Mg fertilizers	75
10 Effects of different sources of Mg on FFB yield	76
11 Yield and leaf analysis result	79
12 P status in oil palm pinnae at Dusun Durian Estate	80
13 Effects of Mg fertilizers on total Mg and K in leaf tissue	80
14 Application of fertilizer at Dusun Durian Estate, Golden Hope Plantations Berhad, Banting, Selangor.....	85
15 Guidelines for the interpretation of correlation coefficient	104
16 Descriptive statistics of N in soil and leaf tissue	107
17 Correlation matrix independent variable of total N in soil	109
18 Relationship between total N with total OC and exchangeable K and Mg in soil	110

19	Relationship between N in soil and leaf tissue	113
20	Geostatistical parameters of N in soil and leaf tissue	115
21	Descriptive statistics of P in soil and leaf tissue	120
22	Correlation matrix independent variable of available P in soil	123
23	Relationship between soil available P with soil pH, exchangeable of Ca and Mg	124
24	Relationship between P in soil and leaf tissue	127
25	Geostatistical parameters of P in soil and leaf tissue	128
26	Descriptive statistics of K in soil and leaf tissue	133
27	Correlation matrix independent variable of exchangeable K in soil	134
28	Relationship between soil exchangeable K with soil pH and exchangeable of Ca and Mg	135
29	Relationship between K in soil and leaf tissue	137
30	Geostatistical parameters of K in soil and leaf tissue	139
31	Descriptive statistics of Ca in soil and leaf tissue	144
32	Correlation matrix independent variable of exchangeable Ca in soil	146
33	Relationship between soil exchangeable Ca with available P and soil pH	147
34	Relationship between Ca in soil and leaf tissue	148
35	Geostatistical parameters of Ca in soil and leaf tissue	149
36	Descriptive statistics of Mg in soil and leaf tissue	154
37	Correlation matrix independent variable of exchangeable Mg in soil	156
38	Relationship between soil exchangeable Mg with total N, available P, exchangeable K, total OC and soil pH	157
39	Relationship between Mg in soil and leaf tissue	158
40	Geostatistical parameters of Mg in soil and leaf tissue	159
41	Descriptive statistics of FFB	164

42	Correlation matrix independent variable of FFB	166
43	Relationship between leaf nutrient and FFB	167
44	Geostatistical parameters of FFB	169
45	Correlation matrix independent variable of DN of SPOT image	171

LIST OF FIGURES

Figure	Pages
1 Elements and system of precision farming	9
2 DGPS Systems	24
3 Tractor Robot Spraying	26
4 Precision Agriculture/Precision Farming Cycle	32
5 Soil Sampler Vehicle	33
6 Typical variogram model	38
7 Kriged maps of total N, available P and exchangeable K in soil	41
8 Typical of Spraying Variable Rate Technology Component	43
9 Variable rate application using a VRT sprayer	43
10 Spectral reflectance curve for various vegetation in Raub	50
11 Biomass classes using temporal images	51
12 Spectral reflectance of soybean leaves damage by <i>Spodoptera litura</i>	52
13 Multi spectral imagery is used to monitor Vineyard health and vigour	53
14 Nitrogen cycle in the soil	65
15 Phosphorous dynamics system in the soil	68
16 The dynamic of K processes in the soil	71
17 Simple equilibrium of Ca and Mg in the soil	73
18 Typical relationship between FFB yield and leafN	77
19 Location of the study site	82
20 The average monthly rainfall from 1990 to 2000 taken at Dusun Durian Estate, Banting	83

21	Clean circle weeding system	84
22	Drainage system in the study area	86
23	Annual production of FFB in the study area	87
24	Method of leaf sample sites collection	88
25	Determination of 17 th frond and leaflets cutting	89
26	Soil and leaf sampling sites	90
27	Weighing of a FFB	91
28	Flowchart of the data analysis	92
29	Pre-sampling process of soil analysis	97
30	Pre-sampling process of leaf tissue analysis	100
31	DGPS measurement in the field	102
32	Transferring field DGPS data into computer	103
33	Location of leaf sample sites	103
34	Relationship between total N and OC at top-layer.....	111
35	Relationship between total N and OC at sub-layer	111
36	Semivariograms of N in soil and leaf tissue	114
37	Spatial distribution of total N in soil	116
38	Spatial distribution of N concentration in leaf tissue	118
39	Distribution of organic P and C in soil	122
40	The inorganic fixation P at various soil pH value	125
41	Effect of pH on the distribution of P ions in soil solution	126
42	Semivariograms of P in soil and leaf tissue	128
43	Spatial distribution of available P in soil	129
44	Spatial distribution of P concentration in leaf tissue	130

45	Relationship between exchangeable K and exchangeable Mg at top-layer	135
46	Semivariograms of K in soil and leaf tissue	138
47	Spatial distribution of exchangeable K in soil	140
48	Spatial distribution of K concentration in leaf tissue	142
49	Semivariograms of Ca in soil and leaf tissue	149
50	Spatial distribution of exchangeable Ca in soil	151
51	Spatial distribution of Ca concentration in leaf tissue	152
52	Semivariograms of Mg in soil and leaf tissue	159
53	Spatial distribution of exchangeable Mg in soil	161
54	Spatial distribution of Mg concentration in leaf tissue	162
55	Semivariogram of FFB	168
56	Spatial distribution of FFB	170
57	SPOT image of the study area	171
58	Unsupervised and Supervised Classification SPOT image of the study area	174
59	pH meter measurement process	198
60	Autoanalyzer measurement process	200
61	Analysis of total N process	202
62	Leaching analysis process	205

CHAPTER I

INTRODUCTION

Background of the Study

The history of oil palm in Malaysia began from the time when oil palm was first introduced into this country in 1890 through the Botanical Gardens, Singapore. However, the first commercial planting was not initiated until 1917 at Tennamaram Estate in Kuala Selangor. Long before its introduction into Malaysia, the oil palm (*Elaeis guineensis* Jacq) was abundantly found in tropical Africa under natural conditions and its kernel and pericarp oil were widely used by the natives (Tang, 1966). In 1968, the area under oil palm was 204,000 ha and increased significantly 10 years later to 850,000 ha (Mielke, 1987). A recent data in 2000 showed that the area under oil palm in Malaysia was 3,463,000 ha (Ministry of Finance Malaysia, 2000), covering about 57.7% of the total area under selected crop plantations in Malaysia (Ministry of Primary Industries Malaysia, 2000).

Production of crude palm oil in Malaysia in 2000 was about 10.7 million tonnes, an increase of 0.9% over the previous year, which made Malaysia the leading producer and exporter country in the world. The latest data showed that during the first eight months in 2000, Malaysian palm oil export was about 5.39 million tonnes compared to 5.48 million tones in 1999 for the same period. The total income of these periods were RM6,576 million and RM9,736 million, respectively (Ministry of Finance Malaysia, 2000; Oil World, 2001). Oil palm

plantation is generally managed conventionally. The conventional technology considered that the whole plantation is homogenous, therefore one set of practice applies to the entire plantation. Such management is inadequate because of the tendency of over treating and under-treating some portions of a field. This may increase input costs, decreased net economic return, and contribute to surface and ground water pollution and waste of energy.

Precision farming (PF) or Precision Agriculture (PA) is a technology available for sustainable agricultural production, which enables farm management based on the small-scale spatial variability of soil and crop variables in the field (Stafford *et al.*, 1996; Haneklaus *et al.*, 1997). The PA aims to achieve a better usage of resources and control mechanism so as to improve production efficiency, reduce input costs and reduce negative environmental impact (Costopoulou and Anagnostou, 1997; Voltz, 1997). Soil, an important component of land should be assessed in terms of their abilities to supply plants need of nutrients. In specifying such plant requirements, it is essential to state them quantitatively and precisely (Tinker and Leigh, 1985). Using precision farming technologies requires better understanding of soil variability in physical and chemical properties. Some of that variations are inherent, but others are the result of the management history of the field (Bocchi *et al.*, 2000). Fundamental question of the concept of precision farming for the farmer is the assessment and appropriate treatment of substantial heterogeneity for site-specific management decision at a reasonable cost (Grenzdorffer, 1997; Van Groenigen *et al.*, 2000). Proper land management leading to sustainable agriculture production system, which are in balance with nature and

the environment, is being guided by farmer's assessment of variation in space and time of soil and crop conditions (Bouma, 1997a).

Problem Statement

In general, oil palm plantations are managed based on conventional technology on large-scale extensive agricultural practices. It appears inevitable to intensify and improve the planting practices by developing new techniques to increase efficiency and productivity to keep ahead of competitors and maintain its reputation as a responsible “green industry”. The conventional technology manages the fields as if they were a homogenous unit, whereas the real condition indicated that land characteristics are very heterogeneous. Conventional crop production methods developed within the last 30-50 years have been linked to the negative effects on the environment, human health and safety, and long term effects on soil fertility (Poudel *et al.*, 2001). Lawrence and William (1990) reported that agricultural production processes generate residuals such as manure, fertilizer, pesticides, and soil particles, which can contaminate both ground and surface water. Numerous health and environmental quality problems have been associated with these contaminants. Spatial variability can occur on a variety of scales, between regions, between fields, or within fields, especially variation in soil properties (Burrough, 1993). Soil chemical properties that vary spatially include nutrients, pH, salinity and organic matter. Soil nutrients are essential for crop growth and the advent of fertilizer technology had made them easily manageable (McBratney and Pringle, 1997). The nutrient balances and the dosage of fertilizers needed to secure

the level of production required and ensure that soil fertility is maintained and preferably improved. Local balances that has sound scientific and economic basis are also required for cropping systems to provide good advice to farmers (Cooke, 1986).

For years, agronomist and soil scientists have encouraged farmers to regularly sample and analyze the soil of their fields. Regular soil sampling is important for developing a successful fertility management program. However, in the past, soil sampling was often overlooked and fertilizer was frequently over-applied in order to guarantee that nutrients levels were adequate. Foliar analysis is carried out to quantify the deficiency of individual nutrients so that fertilizer adjustments can be calculated. Optimum leaf nutrient levels correspond to maximum yield (Foster *et al.*, 1987).

In precision farming, information on soil and plant nutrient status of oil palm is very important and useful for fertilizer recommendation. A PF system matches resource availability to crop capability and then knowledge of soil spatial variability (McBratney and Pringle, 1997). This requires proper information on specific soil conditions, especially soil characteristics and soil behavior. However, soil-specific management will be relevant only if the agronomist is able to deliver accurate site-specific advice (Voltz, 1997). The major problem remains with knowledge of the critical factors and their effects on growth and yield as well as knowledge of important nutrient budget requirements in the oil palm systems. Using global positioning system (GPS) and geographic information system (GIS) technology

would have a quick result in improved availability of information for planning, organization, monitoring and supervision of work. The most useful application will be the integration of agronomic and management information in the establishment of a decision support system (DSS) for oil palm management to assist in site specific decision making for particular blocks (Chew, 1998).

The question is where precision farming can be implemented and how these new technologies for precision planting practices (PPP) can help achieve the purpose for which the plantations were established. Remote sensing image and techniques have to be cost effective, problem oriented, geo-referred and relatively quick, available for a broad range of applications in site-specific management (Grenzdorver, 1997). Remote sensing technology combined with GIS/GPS could be used to obtain information about healthy vegetation and nutrient content in leaf of oil palm plantation. Satellite data such as SPOT can be differentiated in contrast between high and low content of nutrients in plant tissue of oil palm plantation for nitrogen, phosphorous, potassium, calcium and magnesium (Nguyen *et al.*, 1993). Lukman Fadli and Siahaan (1993) reported a good relationship between near infrared (NIR) reflectance with macro nutrient (N, P, K, Ca, Mg, S) status in leaf tissue using SPOT image.

Objective of Study

The general objective of this study is to obtain accurate and proper time information about soil and leaf nutrient variability using GIS and remote sensing